

First-Forbidden Continuum- and Bound-State β^- -Decay Rates of Bare $^{205}\text{Hg}^{80+}$ and $^{207}\text{Tl}^{81+}$ Ions

M. Faber^a, A. N. Ivanov^{a,b}, P. Kienle^{b,c}, E. L. Kryshen^d, M. Pitschmann^a, N. I. Troitskaya^e

^aAtominstytut der Österreichischen Universitäten, Technische Universität Wien,
Wiedner Hauptstrasse 8-10, A-1040 Wien, Österreich

^bStefan Meyer Institut für subatomare Physik Österreichische Akademie der Wissenschaften,
Boltzmannngasse 3, A-1090, Wien, Österreich

^cPhysik Department, Technische Universität München, D-85748 Garching, Germany

^dPetersburg Nuclear Physics Institute, 188300 Gatchina, Orlova roscha 1, Russian Federation and

^ePetersburg Nuclear Physics Institute, 188300 Gatchina, Orlova roscha 1, Russian Federation,

^eState Polytechnic University of St. Petersburg, Polytechnicheskaya 29, 195251, Russian*

(Dated: October 13, 2008)

We calculate the decay rates $\lambda_{\beta_c^-}$ and $\lambda_{\beta_b^-}$ of the continuum- and bound-state β^- -decays for bare $^{205}\text{Hg}^{80+}$ and $^{207}\text{Tl}^{81+}$ ions. For the ratio of the decay rates $R_{b/c} = \lambda_{\beta_b^-}/\lambda_{\beta_c^-}$ we obtain the values $R_{b/c} = 0.161$ and $R_{b/c} = 0.190$ for bare $^{205}\text{Hg}^{80+}$ and $^{207}\text{Tl}^{81+}$ ions, respectively. The theoretical value of the ratio $R_{b/c} = 0.190$ for the decays of $^{207}\text{Tl}^{81+}$ agrees within 1 % of accuracy with the experimental data $R_{b/c}^{\text{exp}} = 0.188(18)$, obtained at GSI. The theoretical ratio $R_{b/c} = 0.161$ for $^{205}\text{Hg}^{80+}$ is about 20 % smaller than the experimental value $R_{b/c}^{\text{exp}} = 0.20(2)$, measured recently at GSI.

PACS: 12.15.Ff, 13.15.+g, 23.40.Bw, 26.65.+t

As has been shown in [1], the standard theory of weak interactions of heavy ions [2], allows to describe well the K-shell electron capture (*EC*) and β^+ decays of the H-like and He-like ions $^{140}\text{Pr}^{58+}$ and $^{140}\text{Pr}^{57+}$, agreeing with the experimental data, obtained at GSI [3], with an accuracy better than 3 %.

In this letter we apply the standard theory of weak interactions of heavy ions [2] and technique developed in [1] to the calculation of the decay rates $\lambda_{\beta_c^-}$ and $\lambda_{\beta_b^-}$ of the decays

$$\begin{aligned} ^{205}\text{Hg}^{80+} &\rightarrow ^{205}\text{Tl}^{81+} + e^- + \tilde{\nu}_e, \\ ^{205}\text{Hg}^{80+} &\rightarrow ^{205}\text{Tl}^{80+} + \tilde{\nu}_e, \\ ^{207}\text{Tl}^{81+} &\rightarrow ^{207}\text{Pb}^{82+} + e^- + \tilde{\nu}_e, \\ ^{207}\text{Tl}^{81+} &\rightarrow ^{207}\text{Pb}^{81+} + \tilde{\nu}_e, \end{aligned} \quad (1)$$

where $^{205}\text{Hg}^{80+}$, $^{207}\text{Pb}^{82+}$ and $^{205}\text{Tl}^{81+}$, $^{207}\text{Tl}^{81+}$ are two pairs of bare ions with quantum numbers $I^P = \frac{1}{2}^-$ and $I^P = \frac{1}{2}^+$, respectively, and $^{205}\text{Tl}^{80+}$ and $^{207}\text{Pb}^{81+}$ are the H-like ions. The continuum- and bound-state β^- -decays in Eq.(1) satisfy the selection rule $\Delta I^P = 0^-$, which corresponds to the selection rule of the first-forbidden β -decays [4]. The bound-state β^- -decay to the K-shell is the time reversed orbital K-shell Electron Capture (*EC*) decay, which we analysed in [1]. The main distinction of the bound state β^- -decay from the *EC*-decay is that the bound electron can be not only on the K-shell in the 1s state but on any other shells in any excited *ns* state, the contribution of which is about 20 %.

A measurement of the continuum- and bound-

state β^- decays of bare $^{207}\text{Tl}^{81+}$ ion was reported by Ohtsubo *et al.* [5]. The experimental value of the ratio of the decay rates $R_{b/c}^{\text{exp}} = 0.188(18)$ agrees within one standard deviation with the theoretical value $R_{b/c}^{\text{th}} = 0.171(1)$ [6], obtained from the theory employing spectra of allowed transitions [5]. Very recently an experimental value $R_{b/c}^{\text{exp}} = 0.20(2)$ of the ratio of the continuum- and bound-state β^- decays of bare $^{205}\text{Hg}^{80+}$ has become known [7]. This has motivated us to carry out a model-independent calculation of the first-forbidden continuum- and bound-state β^- decays of bare $^{205}\text{Hg}^{80+}$ and $^{207}\text{Tl}^{81+}$ ions.

For the calculation of the rates of the β^- -decays we use the Hamilton density operator [1, 2]

$$\begin{aligned} \mathcal{H}_W(x) &= \frac{G_F}{\sqrt{2}} V_{ud} [\bar{\psi}_p(x) \gamma^\mu (1 - g_A \gamma^5) \psi_n(x)] \\ &\times [\bar{\psi}_e(x) \gamma_\mu (1 - \gamma^5) \psi_{\nu_e}(x)] + \text{h.c.} \end{aligned} \quad (2)$$

with standard notations [1, 2, 8]. In our calculations the anti-neutrino is assumed to be a massless Dirac anti-particle as in [1].

The continuum-state β^- -decay

In the rest frame of the mother ion the amplitudes of the continuum-state β^- -decay are defined by

$$\begin{aligned} M_{II_z \rightarrow I' I'_z} &= \\ &= -I' I'_z \langle \tilde{\nu}_e(\vec{k}) e^-(\vec{p}_-) d(\vec{q}) | \mathcal{H}_W(0) | m(\vec{0}) \rangle_{II_z}, \end{aligned} \quad (3)$$

where $II_z = \frac{1}{2}, \pm\frac{1}{2}$ and $I'I'_z = \frac{1}{2}, \pm\frac{1}{2}$ determine the spinorial states of the mother m and daughter d ions. The decay rate $\lambda_{\beta_c^-}$ of the continuum-state β^- -decay is defined by [1]

$$\lambda_{\beta_c^-} = \frac{1}{2M_m} \int \frac{d^3\vec{q}}{(2\pi)^3 2E_d} \frac{d^3\vec{p}_-}{(2\pi)^3 2E_-} \frac{d^3\vec{k}}{(2\pi)^3 2E_{\bar{\nu}_e}} \times (2\pi)^4 \delta^{(4)}(k + p_- + q - k_m) F(Z+1, E_-) \times \frac{1}{2} \sum_{I_z, I'_z, \sigma_-} |M_{II_z \rightarrow I'I'_z}|^2, \quad (4)$$

where $k = (E_{\bar{\nu}_e}, \vec{k})$, $p_- = (E_-, \vec{p}_-)$, $q = (E_d, \vec{q})$ and $k_m = (M_m, \vec{0})$ are 4-momenta of the interacting particles, $F(Z+1, E_-)$ is the Fermi function [9]

$$F(Z+1, E_-) = \left(1 + \frac{1}{2}\gamma\right) \frac{4(2Rp_-)^{2\gamma}}{\Gamma^2(3+2\gamma)} \times e^{+\frac{\pi(Z+1)\alpha E_-}{p_-}} \left| \Gamma\left(1 + \gamma + i \frac{\alpha(Z+1)E_-}{p_-}\right) \right|^2, \quad (5)$$

with $\gamma = \sqrt{1 - ((Z+1)\alpha)^2} - 1$, $p_- = \sqrt{E_-^2 - m_e^2}$, $Z = 80$ and $Z = 81$ for $^{205}\text{Hg}^{80+}$ and $^{207}\text{Tl}^{81+}$, respectively. The summation in Eq.(4) should be carried out over all polarisations of the interacting particles, where $\sigma_- = \pm\frac{1}{2}$ is a polarisation of the electron. The anti-neutrino is polarised parallel to the momentum $\vec{k}$. Following [1], for the amplitudes of the continuum-state β^- -decay we get the expressions

$$\begin{aligned} M_{\frac{1}{2}, +\frac{1}{2} \rightarrow \frac{1}{2}, +\frac{1}{2}} &= \sqrt{2M_m 2E_d} \mathcal{M}_{m \rightarrow d} \\ &\times [\bar{u}_e(\vec{p}_-, \sigma_-) (g_A \gamma^0 - \gamma^3) (1 - \gamma^5) v_{\bar{\nu}_e}(\vec{k}, +\frac{1}{2})], \\ M_{\frac{1}{2}, +\frac{1}{2} \rightarrow \frac{1}{2}, -\frac{1}{2}} &= \sqrt{2M_m 2E_d} \mathcal{M}_{m \rightarrow d} \\ &\times [\bar{u}_e(\vec{p}_-, \sigma_-) (\gamma^1 + i\gamma^2) (1 - \gamma^5) v_{\bar{\nu}_e}(\vec{k}, +\frac{1}{2})], \\ M_{\frac{1}{2}, -\frac{1}{2} \rightarrow \frac{1}{2}, +\frac{1}{2}} &= \sqrt{2M_m 2E_d} \mathcal{M}_{m \rightarrow d} \\ &\times [\bar{u}_e(\vec{p}_-, \sigma_-) (\gamma^1 - i\gamma^2) (1 - \gamma^5) v_{\bar{\nu}_e}(\vec{k}, +\frac{1}{2})], \\ M_{\frac{1}{2}, -\frac{1}{2} \rightarrow \frac{1}{2}, -\frac{1}{2}} &= \sqrt{2M_m 2E_d} \mathcal{M}_{m \rightarrow d} \\ &\times [\bar{u}_e(\vec{p}_-, \sigma_-) (g_A \gamma^0 + \gamma^3) (1 - \gamma^5) v_{\bar{\nu}_e}(\vec{k}, +\frac{1}{2})]. \quad (6) \end{aligned}$$

where $\bar{u}_e$ and $v_{\bar{\nu}_e}$ are Dirac bispinors of the electron and the anti-neutrino, $\mathcal{M}_{m \rightarrow d}$ is the nuclear matrix element defined by

$$\mathcal{M}_{m \rightarrow d} = -\frac{G_F}{\sqrt{2}} V_{ud} \int d^3x \Psi_d^*(\vec{r}) \Psi_m(\vec{r}), \quad (7)$$

where $\Psi_d(\vec{r})$ and $\Psi_m(\vec{r})$ are the wave functions of the daughter and mother nuclei. For the numerical calculations we assume that the product

$\Psi_d^*(\vec{r}) \Psi_m(\vec{r})$ has the Wood-Saxon shape [1]. Substituting the amplitudes Eq.(6) into Eq.(4) and carrying out the summation over polarisations we get

$$\lambda_{\beta_c^-} = (3 + g_A^2) \frac{|\mathcal{M}_{m \rightarrow d}|^2}{\pi^3} f(Q_{\beta_c^-}, Z+1), \quad (8)$$

where the Fermi integral $f(Q_{\beta_c^-}, Z+1)$ is

$$f(Q_{\beta_c^-}, Z+1) = \int_{m_e}^{Q_{\beta_c^-} + m_e} (Q_{\beta_c^-} + m_e - E_-)^2 \times F(Z+1, E_-) \sqrt{E_-^2 - m_e^2} E_- dE_- \quad (9)$$

The Q -values of the continuum-state β^- -decays are equal to $Q_{\beta_c^-} = 1515.734 \text{ keV}$ and $Q_{\beta_c^-} = 1407.471 \text{ keV}$ for $^{205}\text{Hg}^{80+}$ and $^{207}\text{Tl}^{81+}$, respectively [9, 10, 11].

The bound-state β^- -decay

In the bound-state β^- -decay the electron in the final state can be in any bound ns -state. Due to hyperfine interactions the ns -state is splitted into two hyperfine states $(ns)_{F=0}$ and $(ns)_{F=1}$ with a total spin $F = 0$ and $F = 1$, respectively [12].

The experimental value of the hyperfine energy level splitting for the $1s$ -state $\Delta E_{1s}^{\text{exp}} = E_{(1s)_{F=0}} - E_{(1s)_{F=1}} = -3.24409 \pm 0.00029 \text{ eV}$ [13] agrees well with the theoretical one $\Delta E_{1s}^{\text{th}} = -3.275 \text{ eV}$ [12]. The hyperfine splitting $\Delta E_{ns}^{\text{th}}$ of the energy level of the excited ns state is related to $\Delta E_{1s}^{\text{th}}$ as [12]

$$\Delta E_{ns} = \frac{\Delta E_{1s}}{(3+2\gamma)[(n+\gamma)^2 - \gamma(2+\gamma)]^2} \times \left[2(n+\gamma) + \sqrt{(n+\gamma)^2 - \gamma(2+\gamma)} \right]. \quad (10)$$

The decay rate $\lambda_{\beta_b^-}$ of the bound-state β^- -decay into any hyperfine states $(ns)_{F=0}$ and $(ns)_{F=1}$ is defined by

$$\begin{aligned} \lambda_{\beta_b^-} &= \sum_{n=1}^{\infty} \sum_{F=0,1} \lambda_{\beta_b^-}((ns)_F) = \frac{1}{2M_m} \\ &\times \int \frac{d^3\vec{q}}{(2\pi)^3 2E_d} \frac{d^3\vec{k}}{(2\pi)^3 2E_{\bar{\nu}_e}} (2\pi)^4 \delta^{(4)}(q + k - k_m) \\ &\times \frac{1}{2} \sum_{n=1}^{\infty} \sum_{F=0,1} \sum_{I_z, M_F} |M_{II_z \rightarrow F M_F}^{(n)}|^2, \quad (11) \end{aligned}$$

where $k = (E_{\bar{\nu}_e}, \vec{k})$, $q = (E_d, \vec{q})$ and $k_m = (M_m, \vec{0})$ are 4-momenta of the interacting particles. In the rest frame of the mother ion the amplitudes of the bound-state β^- -decay are determined

by

$$M_{II_z \rightarrow FM_F}^{(n)} = -FM_F \langle \tilde{\nu}_e(\vec{k}) d^{(n)}(\vec{q}) | \mathcal{H}_W(0) | (\vec{0}) \rangle_{II_z}, \quad (12)$$

Following [1] for the amplitudes of the transitions $II_z \rightarrow FM_F$ we get the expressions

$$\begin{aligned} M_{\frac{1}{2}, -\frac{1}{2} \rightarrow 0, 0}^{(n)} &= \sqrt{2M_m 2E_d E_{\tilde{\nu}_e}} (3 - g_A) \\ &\times \sqrt{2} \mathcal{M}_{m \rightarrow d} \langle \psi_{ns}^{(Z+1)} \rangle, \\ M_{\frac{1}{2}, +\frac{1}{2} \rightarrow 1, +1}^{(n)} &= \sqrt{2M_m 2E_d E_{\tilde{\nu}_e}} (1 + g_A) \\ &\times 2 \mathcal{M}_{m \rightarrow d} \langle \psi_{ns}^{(Z+1)} \rangle, \\ M_{\frac{1}{2}, -\frac{1}{2} \rightarrow 1, 0}^{(n)} &= \sqrt{2M_m 2E_d(\vec{q}) E_{\tilde{\nu}_e}(\vec{k})} (1 + g_A) \\ &\times \sqrt{2} \mathcal{M}_{m \rightarrow d} \langle \psi_{ns}^{(Z+1)} \rangle, \end{aligned} \quad (13)$$

where $\langle \psi_{ns}^{(Z+1)} \rangle$ is the wave function of the bound electron in the ns -state, averaged over the nuclear density [1]. The wave function $\psi_{ns}^{(Z+1)}$ is the solution of the Dirac equation [15, 16]. Substituting the amplitudes Eq.(13) into Eq.(11) we get the decay rate

$$\begin{aligned} \lambda_{\beta_b^-} &= (3 - g_A)^2 |\mathcal{M}_{m \rightarrow d}|^2 \sum_{n=1}^{\infty} |\langle \psi_{ns}^{(Z+1)} \rangle|^2 \frac{Q_{ns}^2}{2\pi} \\ &+ 3(1 + g_A)^2 |\mathcal{M}_{m \rightarrow d}|^2 \sum_{n=1}^{\infty} |\langle \psi_{ns}^{(Z+1)} \rangle|^2 \frac{Q_{ns}^2}{2\pi} \\ &= 2(3 + g_A^2) |\mathcal{M}_{m \rightarrow d}|^2 \sum_{n=1}^{\infty} |\langle \psi_{ns}^{(Z+1)} \rangle|^2 \frac{Q_{ns}^2}{\pi}, \end{aligned} \quad (14)$$

where Q_{ns} is the Q -value of the bound-state β^- -decays into the bound ns -state [14].

The ratio of the continuum- and bound-state β^- -decays

The ratio of the continuum- and bound-state β^- -decay rates is

$$R_{\beta_b^-/\beta_c^-} = \sum_{n=1}^{\infty} \frac{2\pi^2 Q_{ns}^2 |\langle \psi_{ns}^{(Z+1)} \rangle|^2}{f(Q_{\beta_c^-}, Z+1)}. \quad (15)$$

The numerical values of the ratio of the β^- -decays of $^{205}\text{Hg}^{80+}$ and $^{207}\text{Tl}^{81+}$ are

$$\begin{aligned} ^{205}\text{Hg}^{80+} : \quad R_{\beta_b^-/\beta_c^-}^{\text{th}} &= 0.161, \\ ^{207}\text{Tl}^{81+} : \quad R_{\beta_b^-/\beta_c^-}^{\text{th}} &= 0.190. \end{aligned} \quad (16)$$

These results, obtained for $R = 1.1 \times A^{1/3}$ fm [1, 13], are stable under reasonable variations of R as it has been observed for the EC and β^+ decays of the H-like $^{140}\text{Pr}^{58+}$ and He-like $^{140}\text{Pr}^{57+}$ ions in [1].

Concluding discussion

Using the standard theory of weak interactions of heavy ions we have calculated the decay rates of the continuum- and bound-state β^- -decays of bare $^{205}\text{Hg}^{80+}$ and $^{207}\text{Tl}^{81+}$ ions. These are first -forbidden β^- -decays [4]. Our result for the ratio of the β^- -decays of bare $^{207}\text{Tl}^{81+}$ ion $R_{\beta_b^-/\beta_c^-}^{\text{th}} = 0.190$ agrees with the experimental value $R_{\beta_b^-/\beta_c^-}^{\text{exp}} = 0.188(18)$ within an accuracy of about 1%. This is much better than the theoretical value $R_{\beta_b^-/\beta_c^-}^{\text{th}} = 0.171(1)$, obtained by Takahashi and Yokoi [6].

Our result for the ratio of the β^- -decay rates of bare $^{205}\text{Hg}^{80+}$ ions deviates from the experimental value $R_{\beta_b^-/\beta_c^-}^{\text{exp}} = 0.20(2)$ by about 20%. Since our agreement with the experimental value of $R_{\beta_b^-/\beta_c^-}^{\text{exp}} = 0.188(18)$ for bare $^{207}\text{Tl}^{81+}$ ions is about of 1%, we can argue that the experimental value for the ratio of the β^- -decay rates of bare $^{205}\text{Hg}^{80+}$ ions maybe actually too high. Our assertion is based on the following. The dependence of the ratio $R_{\beta_b^-/\beta_c^-}$ on the electric charge is rather weak. Indeed, for $Z = 81$ instead of $Z = 80$ the ratio of the β^- -decays $R_{\beta_b^-/\beta_c^-}$ of bare $^{205}\text{Hg}^{80+}$ ions changes to the value $R_{\beta_b^-/\beta_c^-} = 0.167$. Thus, the main distinction is due to different Q -values of the continuum- and bound-state β^- -decays. The Q -values $Q_{\beta_c^-} = 1515.734$ keV and $Q_{\beta_b^-} = 1407.471$ keV of continuum-state β^- -decays result in the Fermi integrals $f(Q_{\beta_c^-}, Z+1) = 22.119$ MeV⁵ and $f(Q_{\beta_b^-}, Z+1) = 17.747$ MeV⁵ for $^{205}\text{Hg}^{80+}$ and $^{207}\text{Tl}^{81+}$, respectively. On the other hand the bound-state β^- -decay rates scale with Q_{ns}^2 , where $Q_{1s} = 1614.557$ keV and $Q_{1s} = 1509.053$ keV for $^{205}\text{Hg}^{80+}$ and $^{207}\text{Tl}^{81+}$, respectively. This implies that the ratio of the continuum- and bound-state β^- -decay rates for bare $^{205}\text{Hg}^{80+}$ ions should be as minimum 1.1 times smaller relative to the ratio of the continuum- and bound-state β^- -decay rates for bare $^{207}\text{Tl}^{81+}$ ions. This gives $R_{\beta_b^-/\beta_c^-} \sim 0.171$ for bare $^{205}\text{Hg}^{80+}$ ions at the nuclear radius $R = 1.1 \times A^{1/3}$. A reduction of the value $R_{\beta_b^-/\beta_c^-} \sim 0.171$ to $R_{\beta_b^-/\beta_c^-} = 0.161$ is caused by the effective densities of electrons in the ns -states for different $(Z+1)$ values of electric charges of the ions $^{205}\text{Tl}^{80+}$ and $^{207}\text{Pb}^{81+}$ in the final state of the bound-state β^- -decays.

-
- * Electronic address: ivanov@kph.tuwien.ac.at
- [1] A. N. Ivanov, M. Faber, R. Reda, P. Kienle, Phys. Rev. C **78**, 025503, arXiv: 0711.3184 [nucl-th].
- [2] W. Bambynek *et al.*, Rev. Mod. Phys. **49**, 77 (1977).
- [3] Yu. A. Litvinov *et al.* (the GSI Collaboration), Phys. Rev. Lett. **99**, 262501 (2008).
- [4] E. J. Konopinski, in *The theory of beta radioactivity*, Oxford, At the Clarendon Press, 1966.
- [5] T. Ohtsubo *et al.* (the GSI Collaboration), Phys. Rev. Letters, **95**, 052501 (2004).
- [6] K. Takahashi and K. Yokoi, Nucl. Phys. A **404**, 578 (1983); Atomic Data and Nuclear Data Tables **36**, 375 (1987).
- [7] J. Kurcewicz *et al.* (the GSI Collaboration), (to be published)
- [8] W.-M. Yao *et al.*, J. Phys. G **33**, 1(2006).
- [9] A. N. Ivanov, E. L. Kryshen, M. Pitschmann, P. Kienle, Phys. Rev. Letters (in press), arXiv: 0806.2543 [nucl-th].
- [10] The Q -value of the continuum-state β^- -decays we calculate as: $Q_{\beta_c^-} = (\Delta_m - \Delta_d) + (\epsilon_{TF}^{(Z+1)} - \epsilon_{TF}^{(Z)}) = (-22304 + 23835) + (-521.826 + 506.560) \text{ keV} = 1515.734 \text{ keV}$ for $^{205}\text{Hg}^{80+} \rightarrow$
- $^{205}\text{Tl}^{81+} + e^- + \tilde{\nu}_e$ and $Q_{\beta_c^-} = (\Delta_m - \Delta_d) + (\epsilon_{TF}^{(Z+1)} - \epsilon_{TF}^{(Z)}) = (-21044 + 22467) + (-537.355 + 521.826) \text{ keV} = 1407.471 \text{ keV}$ for $^{207}\text{Tl}^{81+} \rightarrow ^{207}\text{Pb}^{82+} + e^- + \tilde{\nu}_e$, where $\epsilon_{TF}^{(Z)}$ = $-14.33 Z^{2.39} \text{ eV}$ is the binding energy of electrons in the atom with Z electrons, defined in the Thomas-Fermi model (see also [11]).
- [11] R. B. Firestone *et al.*, in *Table of Isotopes*, Eighth Edition Vol.II, A Wiley-Interscience Publication, John Wiley & Sons, Inc, New York, 1996.
- [12] V. M. Shabaev, J. Phys. B: At. Mol. Opt. Phys. **27**, 5825 (1994); V. M. Shabaev *et al.*, Phys. Rev. A **56**, 252 (1997); M. Tomaselli *et al.*, Phys. Rev. A **65**, 022502 (2002).
- [13] P. Beiersdorfer *et al.*, Phys. Rev. A **64**, 032506 (2001); Nucl. Instr. and Meth. Phys. Res. B **205**, 62 (2003).
- [14] The Q -value of the bound-state β^- -decay is defined by $Q_{ns} = Q_{\beta_c^-} + m_e(1 - \frac{n+\gamma}{\sqrt{n^2+2(n-1)\gamma}})$.
- [15] M. E. Rose, in *Relativistische Elektronentheorie*, Band II, pp.218–219, 1971.
- [16] L. D. Landau und E. M. Lifschitz, im *Relativistische Quantentheorie*, Band IVa von W. B. Berestetzki, E. M. Lifschitz und L. P. Pitajewski, Akademie-Verlag, Berlin, 1971.